

Grazing and Haying Restrictions for Herbicides Used in Pasture and Rangeland.

| Herbicide ¹ | Lactating dairy animals | | All animals except lactating dairy animals | | |
|---|--|--------------------|--|--------------------|--------------------------|
| | Before grazing | Before hay harvest | Before Grazing ⁹ | Before hay harvest | Removal before slaughter |
| | ----- days after application ----- | | ----- days after application ----- | | |
| Amber | 0 | 30 | 0 | 30 | 0 |
| Cimarron Max* | 7 | - | 0 | 37 | 30 |
| Cimarron Xtra* | 0 | 0 | 0 | 0 | 0 |
| Crossbow | Next season | Next season | 0 ² | 14 | 3 |
| Curtail* | 14 | 7 | 0 | 7 | 7 ³ |
| Dicamba ¹ | | | | | |
| Up to 1 pt | 7 | 37 | 0 | 0 | 30 |
| Up to 2 pt | 21 | 51 | 0 | 0 | 30 |
| Up to 4 pt | 40 | 70 | 0 | 0 | 30 |
| Up to 16 pt | 60 | 90 | 0 | 0 | 30 |
| Escort* | 0 | 0 | 0 | 0 | 0 |
| Facet L | No grazing restrictions. Allow 7 days before haying. | | | | |
| Glyphosate ¹ | | | | | |
| Pre/Renovation - < 2.25 lb ae/A | No restriction | No restriction | No restriction | No restriction | No restriction |
| - > 2.25 lb ae/A | 8 weeks | 8 weeks | 8 weeks | 8 weeks | 8 weeks |
| Spot Spray - Bayer ⁶ | 7 | 7 | 7 | 7 | 7 |
| - Generics | 14 | 14 | 14 | 14 | 14 |
| Broadcast (8-11 fl oz) | No restriction | No restriction | No restriction | No restriction | No Restriction |
| Grazone P+D | 7 | 30 | 0 | 30 | 3 |
| Landmaster BW* ⁵ | 7 | 30 | 0 | 30 | 3 |
| Method | Do not graze or feed | | Do not graze or feed | | - |
| Milestone | 0 | 0 | 0 | 0 | 0 |
| paraquat* ⁴ | 1 month | 1 month | 1 month | 1 month | 0 |
| Perspective | Do not graze or feed | | Do not graze or feed | | - |
| X Plateau | 0 | 7 | 0 | 7 | 0 |
| Rave | 7 | 0 | 0 | 0 | 0 |
| Spike ⁷ | 0 | 1 year | 0 | 1 year | 0 |
| Stinger* | 0 | 0 | 0 | 0 | 0 |
| Telar* (<0.3 oz/A) | 0 | 0 | 0 | 0 | 0 |
| X Tordon 22K ⁸ | 14 | 1 qt =14, <1qt=0 | 0 | 1 qt =14, <1qt=0 | 3 |
| X 2,4-D/MCPA ¹ (labels vary) | 7-14 | 7-30 | 0-7 | 7-30 | 3-7 |

*Or generic equivalent.

¹These are from Roundup Powermax label. Check specific label containing the same active ingredients for restrictions and uses.

²One year if more than 1.5 gallons/A is used.

³Withdrawal not needed if 2 weeks or more of time elapsed since application.

⁴Restrictions based on degree of new seedlings established before grazing. Suggested at least 6 inches of grass or legume growth.

⁵No restrictions if 10% or less of the area is treated.

⁶Do not treat more than one-tenth of any given acre at one time with spot or wiper application. Remove livestock before application.

⁷If no more than 20 lb/A is used.

⁸Remove livestock to untreated grass pastures for 7 days before transferring livestock to broadleaf or pasture areas. Removal before slaughter statement only applies to animals grazing treated forage for 2 weeks immediately after application.

⁹Some labels do not specify a grazing restriction.